

加納航治
2014年1月末日までの全論文

1	Photochemical reduction of nitrobenzene and its reduction intermediates. IV. Photochemical reduction of azoxybenzene and related compounds. Hashimoto, S.; Kano, K.; Sunamoto, J., Kogyo Kagaku Zasshi 1968, 71 (6), 864-8.
2	Photochemical reduction of nitrobenzene and its reduction intermediates. III. Photochemical reduction of nitrobenzene. Hashimoto, S.; Sunamoto, J.; Fujii, H.; Kano, K., Bull. Chem. Soc. Jpn. 1968, 41 (5), 1249-51.
3	Photochemical direct amination of aromatics with ammonia. Hashimoto, S.; Sunamoto, J.; Kano, K.; Noritsugu, H., Doshisha Daigaku Rikogaku Kenkyu Hokoku. Sci. Eng. Rev. Doshisha Univ. 1968, 9 (1), 65-8.
4	Photochemical reduction of 3,6-diphenyl-s-tetrazine. Hashimoto, S.; Kano, K., Doshisha Daigaku Rikogaku Kenkyu Hokoku 1969, 10 (2), 196-203.
5	Photochemical reduction of nitrobenzene and its reduction intermediates. VI. Photochemical reduction of hydroxyazobenzenes. Hashimoto, S.; Kano, K., Kogyo Kagaku Zasshi 1969, 72 (1), 191-4.
6	Photochemical reduction of nitrobenzene and its reduction intermediates. V. Photochemical reduction of α -nitronaphthalene. Hashimoto, S.; Kano, K., Kogyo Kagaku Zasshi 1969, 72 (1), 188-91.
7	Reduction of azobenzene by hydrochloric acid-isopropyl alcohol. Hashimoto, S.; Kano, K.; Nomura, M., Kogyo Kagaku Zasshi 1969, 72 (9), 2157-8.
8	Photochemical reduction of nitrobenzene and its reduction intermediates. VII. Photochemical reduction of 4-nitropyridine. Preparation of 4,4'-azopyridine. Hashimoto, S.; Kano, K.; Ueda, K., Tetrahedron Lett. 1969, (32), 2733-6.
9	Photochemical reduction of nitrobenzene and its reduction intermediates. VIII. Photochemical reduction of para-substituted nitrobenzenes in isopropanol. Hashimoto, S.; Kano, K., Tetrahedron Lett. 1970, (40), 3509-12.
10	Photochemical reduction of aromatic nitro compounds. Hashimoto, S.; Kano, K., Kagaku (Kyoto) 1971, 26 (12), 1210-18.
11	Photochemical reduction of nitrobenzene and its reduction intermediates. IX. Photochemical reduction of 4-nitropyridine in a hydrochloric acid-isopropyl alcohol solution. Hashimoto, S.; Kano, K.; Ueda, K., Bull. Chem. Soc. Jpn. 1971, 44 (4), 1102-6.
12	Photochemical reduction of nitrobenzene and its reduction intermediates. XI. Photochemical reduction of azobenzene in isopropyl alcohol. Hashimoto, S.; Kano, K., Bull. Chem. Soc. Jpn. 1972, 45 (3), 852-5.
13	Photochemical reduction of nitrobenzene and its reduction intermediates. X. Photochemical reduction of the monosubstituted nitrobenzenes in 2-propanol. Hashimoto, S.; Kano, K., Bull. Chem. Soc. Jpn. 1972, 45 (2), 549-53.
14	Photochemical reaction of p-benzoquinone in water. Hashimoto, S.; Kano, K.; Okamoto, H., Bull. Chem. Soc. Jpn. 1972, 45 (3), 966.
15	Photochemical reaction of 4-nitropyridine in the hydrochloric acid-2-propanol solutions under an oxygen atmosphere. Hashimoto, S.; Kano, K.; Takada, I., Nippon Kagaku Kaishi 1972, (9), 1690-4.
16	Photochemical rearrangement of benzyl phenyl ether. Hashimoto, S.; Nomura, M.; Kano, K., Nippon Kagaku Kaishi 1972, (1), 92-5.
17	Excited state 4-nitropyridine 1-oxide photochemistry in 2-propanol. Hashimoto, S.; Kano, K.; Takeda, I., Doshisha Daigaku Rikogaku Kenkyu Hokoku 1972, 12 (3), 214-20.
18	Photochemical reaction of the p-benzoquinones in a micellar system. Kano, K.; Matsuo, T., Chem. Lett. 1973, (10), 1127-32.
19	Macrocyclic enzyme-model system. Evidence for incorporation of a substrate into a simple cyclic cavity. Murakami, Y.; Sunamoto, J.; Kano, K., Chem. Lett. 1973, No. 3, 223-6.
20	Photochemistry in micellar system. I. Stabilization of the radical ions of anthraquinonesulfonates. Kano, K.; Matsuo, T., Bull. Chem. Soc. Jpn. 1974, 47 (11), 2836-42.
21	Photochemistry of sodium 1,2-naphthoquinone-4-sulfonate in water. Kano, K.; Matsuo, T., Tetrahedron Lett. 1974, (49/50), 4323-6.
22	Radical anion of sodium 9,10-anthraquinone-2-sulfonate stabilized by binding with micelles. Kano, K.; Matsuo, T., Chem. Lett. 1974, (1), 11-14.
23	Macrocyclic enzyme model system. Acyl transfer from p-nitrophenyl carboxylates to paracyclophane oxime. Murakami, Y.; Sunamoto, J.; Kano, K., Bull. Chem. Soc. Jpn. 1974, 47 (5),
24	Photochemical reduction of pyridinium and pyridine compounds in the presence of diethylamine. Kano, K.; Matsuo, T., Tetrahedron Lett. 1975, (16), 1389-92.

加納航治
2014年1月末日までの全論文

25	Electron transfer from photoreduced products of NAD ⁺ and its analogs to hemin in aqueous solutions. Kano, K.; Shibata, T.; Kajiyara, M.; Matsuo, T., <i>Tetrahedron Lett.</i> 1975, (43), 3693–6.
26	Photochemistry in micellar systems. II. Photochemical reduction of β -arylquinonesulfonates in the presence of cationic surfactants. Kano, K.; Takada, Y.; Matsuo, T., <i>Bull. Chem. Soc. Jpn.</i> 1975, 48 (11), 3215–19.
27	Photoinduced one-electron reduction of 1-benzyl-3-carbamoylpyridinium chloride and 3,5-bis(ethoxycarbonyl)-2,6-dimethylpyridine. Kano, K.; Matsuo, T., <i>Bull. Chem. Soc. Jpn.</i> 1976, 49 (11), 3269–73.
28	Enhanced uptake of drugs in liposomes: use of labile vitamin B ₁₂ complexes of 6-mercaptapurine and 8-azaguanine. Kano, K.; Fendler, J. H., <i>Life Sci.</i> 1977, 20 (10), 1729–34.
29	Pyranine as a sensitive pH probe for liposome interiors and surfaces. pH gradients across phospholipid vesicles. Kano, K.; Fendler, J. H., <i>Biochim. Biophys. Acta, Biomembr.</i> 1978, 509 (2),
30	Interaction of vitamin B12a with 8-azaguanine and 6-mercaptapurine: kinetic and thermodynamic characterizations. Kano, K.; Nome, F.; Fendler, J. H., <i>J. Chem. Soc., Dalton Trans.</i> 1978, (10),
31	Zinc tetraphenylporphyrin-sensitized photoreduction of anthraquinonesulfonate in aqueous micellar solutions. Kano, K.; Takuma, K.; Ikeda, T.; Nakajima, D.; Tsutsui, Y.; Matsuo, T., <i>Photochem. Photobiol.</i> 1978, 27 (6), 695–701.
32	Dynamic fluorescence investigations of the effect of osmotic shocks on the microenvironments of charged and uncharged dipalmitoyl-D,L- α -phosphatidylcholine liposomes. Kano, K.; Fendler, J. H., <i>Chem. Phys. Lipids</i> 1979, 23 (3), 189–200.
33	Characterization of surfactant vesicles as membrane mimetic agents. 2. Temperature-dependent changes of the turbidity, viscosity, fluorescence polarization of 2-methylantracene, and positron annihilation in sonicated dioctadecyldimethylammonium chloride. Kano, K.; Romero, A.; Djermouni, B.; Ache, H. J.; Fendler, J. H., <i>J. Am. Chem. Soc.</i> 1979, 101 (15), 4030–7.
34	Energy transfer in artificial membrane systems. Singlet-singlet energy transfer from alloxazines to isoalloxazines in dipalmitoyl phosphatidylcholine liposomes and dialkylammonium chloride vesicles. Aso, Y.; Kano, K.; Matsuo, T., <i>Biochim. Biophys. Acta, Biomembr.</i> 1980, 599 (2), 403–16.
35	Fluorescence quenching of pyrene and pyrenedecanoic acid by various kinds of N,N-dialkylanilines in dipalmitoylphosphatidylcholine liposomes. Kano, K.; Kawazumi, H.; Ogawa, T.; Sunamoto, J., <i>Chem. Phys. Lett.</i> 1980, 74 (3), 511–14.
36	Transmission of photoexcitation energy in biomembrane-like systems. Kano, K.; Matsuo, T., <i>Kagaku (Kyoto)</i> 1980, 35 (2), 120–9.
37	Three component complexes of cycloheptaamylose. Fluorescence quenching of pyrenes and naphthalenes in aqueous media. Kano, K.; Takenoshita, I.; Ogawa, T., <i>Chem. Lett.</i> 1980, (8), 1035–8.
38	Photoresponsive membranes. Regulation of membrane properties by photoreversible cis-trans isomerization of azobenzenes. Kano, K.; Tanaka, Y.; Ogawa, T.; Shimoura, M.; Okahata, Y.; Kunitake, T., <i>Chem. Lett.</i> 1980, (4), 421–4.
39	A study of intervesicle exchange of fluorescent probes in phospholipid bilayer membranes. Kano, K.; Yamaguchi, T.; Matsuo, T., <i>J. Phys. Chem.</i> 1980, 84 (1), 72–6.
40	Energy transfer in micellar system. Fluorescence investigation of energy transfer from dialkylalloxazine to dye cations in the presence of anionic micelles. Matsuo, T.; Aso, Y.; Kano, K., <i>Ber. Bunsenges. Phys. Chem.</i> 1980, 84 (2), 146–52.
41	Unique Properties of Chromophore-Containing Bilayer Aggregates: Enhanced Chirality and Photochemically Induced Morphological Change. Kunitake, T.; Nakashima, N.; Shimomura, M.; Okahata, Y.; Kano, K.; Ogawa, T. <i>J. Am. Chem. Soc.</i> 1980, 102, 6642–6644
42	Liposome Fusion as Monitored by a Fluorescence Quenching Technique. Sunamoto, J.; Iwamoto, K.; Kondo, H.; Kano, K.; Ogawa, T., <i>Chem. Lett.</i> 1981, 15–18.
43	Photo-induced redox reactions in micelles and related systems. Matsuo, T.; Kano, K.; Takuma, K.; Shuto, Y.; Tsutsui, Y.; Nishijima, T., <i>Water Met. Cations Biol. Syst., [Proc. Symp.]</i> 1980, 83–90.
44	Solvent effects on fluorescence of the intramolecular heteroexcimer system and its use for investigating polarity of the microscopic environment. Kano, K.; Goto, H.; Ogawa, T., <i>Chem. Lett.</i> 1981, (5), 653–6.
45	Water Permeation Controlled by trans-cis Isomerization of Photoresponsive Polymer/Artificial Lipid Composite Membrane. Niwa, O.; Kumano, A.; Kajiyama, T.; Takayanagi, M.; Kunitake, T.; Kano, K., <i>Rep. Progress Polym. Phys. Jpn.</i> 1981, 24, 741–743.
46	Steady-state and time-resolved fluorescence studies on energy transfer in anionic surfactant membranes. Kano, K.; Kawazumi, H.; Ogawa, T., <i>J. Phys. Chem.</i> 1981, 85 (20), 2998–3003.

加納航治
2014年1月末日までの全論文

47	Fluorescence quenching in liposomal membranes. Exciplex as a probe for investigating artificial lipid membrane properties. Kano, K.; Kawazumi, H.; Ogawa, T.; Sunamoto, J., <i>J. Phys. Chem.</i> 1981, 85 (15), 2204–9.
48	Photoresponsive artificial membrane. Regulation of membrane permeability of liposomal membrane by photoreversible cis–trans isomerization of azobenzenes. Kano, K.; Tanaka, Y.; Ogawa, T.; Shimomura, M.; Kunitake, T., <i>Photochem. Photobiol.</i> 1981, 34 (3), 323–9.
49	Energy conversion of light by means of chemical reactions at the interface between water and surfactant molecular assemblies. Matsuo, T.; Kano, K.; Nagamura, T.; Sakamoto, T.; Takuma, K.; Itoh, K.; Nishijima, T.; Takayanagi, T., <i>Kenkyu Hokoku – Asahi Garasu Kogyo Gijutsu Shoreikai</i> 1981,
50	Highly sensitive laser fluorimetry of europium(III) with 1,1,1–trifluoro–4–(2–thienyl)–2,4–butanedione. Yamada, S.; Miyoshi, F.; Kano, K.; Ogawa, T., <i>Anal. Chim. Acta</i> 1981, 127, 195–8.
51	Is it possible to determine microscopic viscosities of micelle and membrane interiors? Kano, K., <i>Bunseki</i> 1982, (12), 931–3.
52	γ -Cyclodextrin–enhanced excimer fluorescence of pyrene and effect of n–butyl alcohol. Kano, K.; Takenoshita, I.; Ogawa, T., <i>Chem. Lett.</i> 1982, (3), 321–4.
53	Fluorescence quenching of pyrene and naphthalene in aqueous cyclodextrin solutions. Evidence of three–component complex formation. Kano, K.; Takenoshita, I.; Ogawa, T., <i>J. Phys. Chem.</i> 1982, 86 (10), 1833–8.
54	Laser two–photon ionization technique for a highly sensitive detection of pyrene. Yamada, S.; Kano, K.; Ogawa, T., <i>Bunseki Kagaku</i> 1982, 31 (7), E247–E250.
55	Time discrimination in the laser fluorimetry and ultratrace determination of europium(III) and samarium(III) with 4,4,4–trifluoro–1–(2–thienyl)–1,3–butanedione. Yamada, S.; Kano, K.; Ogawa, T., <i>Anal. Chim. Acta</i> 1982, 134, 21–9.
56	Ion Permeation Control by Photoresponsive Substance Included in Polymer Composite Membrane. Niwa, O.; Kumano, A.; Kajiyama, T.; Takayanagi, M.; Kunitake, T.; Ksano, K.; Shinkai, S., <i>Rep. Progress Polym. Phys. Jpn.</i> 1982, 25, 797–800.
57	Stimulus and response in synthetic membranes. Kano, K., <i>Kagaku, Zokan (Kyoto)</i> 1983, (98), 73–86.
58	Freeze–thaw effect on solubilization of a hydrophobic fluorescent probe in surfactant micelles. Kano, K.; Ishibashi, T.; Ogawa, T., <i>J. Phys. Chem.</i> 1983, 87 (16), 3010–12.
59	Evidence for stacking of cationic porphyrin in aqueous solution. Kano, K.; Miyake, T.; Uomoto, K.; Sato, T.; Ogawa, T.; Hashimoto, S., <i>Chem. Lett.</i> 1983, (12), 1867–70.
60	Fluorescence quenching of water–soluble porphyrins. A novel fluorescence quenching of anionic porphyrin by anionic anthraquinone. Kano, K.; Sato, T.; Yamada, S.; Ogawa, T., <i>J. Phys. Chem.</i> 1983, 87 (4), 566–9.
61	Photoinduced ion permeation through a ternary composite membrane composed of polymer/liquid crystal/azobenzene–bridged crown ether. Kumano, A.; Niwa, O.; Kajiyama, T.; Takayanagi, M.; Kano, K.; Shinkai, S., <i>Chem. Lett.</i> 1983, (8), 1327–30.
62	Control of characteristics of synthetic bilayer membranes via specific interactions with ions. Kunitake, T.; Shimomura, M.; Iida, K.; Okahata, Y.; Kano, K.; Ogawa, T., <i>Nippon Kagaku Kaishi</i> 1983,
63	Laser two–photon ionization for determination of aromatic molecules in solution. Yamada, S.; Hino, A.; Kano, K.; Ogawa, T., <i>Anal. Chem.</i> 1983, 55 (12), 1914–17.
64	Fluorescence quenching of 5,10,15,20–tetra(p–tolyl)porphine and its zinc complex by quinones. Charge–transfer interaction and transient effect. Yamada, S.; Sato, T.; Kano, K.; Ogawa, T., <i>Photochem. Photobiol.</i> 1983, 37 (3), 257–62.
65	Fluorescence in biomimetic membrane systems. Kano, K., <i>Maku</i> 1984, 9 (3), 161–8.
66	Three–component complexes of cyclodextrins. Exciplex formation in cyclodextrin cavity. Kano, K.; Hashimoto, S.; Imai, A.; Ogawa, T., <i>J. Inclusion Phenom.</i> 1984, 2 (3–4), 737–46.
67	Freeze–thaw effect on the solubilization of hydrophobic compounds in micelles and artificial bilayer membranes. Kano, K.; Ueno, Y.; Umakoshi, K.; Hashimoto, S.; Ishibashi, T.; Ogawa, T., <i>J. Phys. Chem.</i> 1984, 88 (21), 5087–92.
68	Fluorescence studies on morphological change of oil–in–water microemulsions upon dilution with water. Kano, K.; Yamaguchi, T.; Ogawa, T., <i>J. Phys. Chem.</i> 1984, 88 (4), 793–6.
69	Solute–solvent interaction in perfluorohexane. Kano, K.; Yamamori, S.; Yanagimoto, M.; Hashimoto, S., <i>Chem. Lett.</i> 1984, (3), 379–82.
70	Photoresponsive permeation characteristics of a ternary composite membrane of polymer/artificial lipid/azobenzene derivative. Kumano, A.; Niwa, O.; Kajiyama, T.; Takayanagi, M.; Kunitake, T.; Kano, K., <i>Polym. J. (Tokyo)</i> 1984, 16 (6), 461–70.
71	Molecular complexes of anionic porphyrin and anionic aromatics. Sato, T.; Ogawa, T.; Kano, K., <i>J. Phys. Chem.</i> 1984, 88 (16), 3678–82.

加納航治
2014年1月末日までの全論文

72	A chiral pyrene excimer in γ -cyclodextrin cavity. Kano, K.; Matsumoto, H.; Hashimoto, S.; Sisido, M.; Imanishi, Y., <i>J. Am. Chem. Soc.</i> 1985, 107 (21), 6117-18.
73	Fluorescence studies on the characterization and solubilizing abilities of sodium dodecyl sulfate, hexadecyltrimethylammonium chloride, and Triton X-100 micelles. Kano, K.; Ueno, Y.; Hashimoto, S., <i>J. Phys. Chem.</i> 1985, 89 (14), 3161-6.
74	Nucleophilic addition of primary alcohols to N-methylacridinium cation. Kano, K.; Zhou, B.; Hashimoto, S., <i>Chem. Lett.</i> 1985, (6), 791-2.
75	Fluorescence study on characterization of liquid domains formed in a frozen acetone-water mixture. Kano, K.; Zhou, B.; Hashimoto, S., <i>J. Phys. Chem.</i> 1985, 89 (17), 3748-52.
76	Effects of aeration, freezing and thawing on self-aggregation of naphthalene in aqueous β -cyclodextrin solutions. Kano, K.; Zhou, B.; Sakaguchi, M.; Matsumoto, H.; Hashimoto, S., <i>Sci. Eng. Rev. Doshisha Univ.</i> 1985, 25 (4), 253-9.
77	Fluorescence studies on solubilizing abilities of micelles and liposomes. Kano, K.; Ueno, Y.; Taniguchi, M.; Ueda, T.; Hashimoto, S., <i>Kenkyu Hokoku - Asahi Garasu Kogyo Gijutsu Shoreikai</i>
78	Exciplexes in perfluorohydrocarbons. Charge-transfer interaction between exciplexes and perfluorohexane. Kano, K.; Yanagimoto, M.; Hashimoto, S., <i>Bull. Chem. Soc. Jpn.</i> 1986, 59 (11),
79	Fluorescence quenching of perfluoronaphthalene by triethylamine. Kano, K.; Yanagimoto, M.; Uraki, H.; Zhou, B.; Hashimoto, S., <i>Bull. Chem. Soc. Jpn.</i> 1986, 59 (4), 993-6.
80	New Fluorescence from Molecular Aggregates of 10-(1-Pyrenyl)decanoic Acid. Itaya, A.; Kawamura, T.; Masuhara, H.; Taniguchi, Y.; Mitsuya, M.; Uraki, H.; Kano, K.; Hashimoto, S., <i>Chem. Lett.</i> 1986, 1541-1544.
81	Fluorescence quenching of a cationic porphyrin by cationic and anionic aromatics: formation of ground-state complexes. Kano, K.; Nakajima, T.; Hashimoto, S., <i>J. Phys. Chem.</i> 1987, 91 (27), 6614-
82	Self aggregation of cationic porphyrin in water. Kano, K.; Nakajima, T.; Takei, M.; Hashimoto, S., <i>Bull. Chem. Soc. Jpn.</i> 1987, 60 (4), 1281-7.
83	Various self assemblies of 10-(1-pyrenyl)decanoic acid. Kano, K.; Uraki, H.; Hashimoto, S., <i>Nippon Kagaku Kaishi</i> 1987, (3), 556-62.
84	The photoreduction mechanism of 10-methylacridinium chloride in methanol. Formation of 9,10-dihydro-9-methoxy-10-methylacridine and hydride transfer. Kano, K.; Zhou, B.; Hashimoto, S., <i>Bull. Chem. Soc. Jpn.</i> 1987, 60 (3), 1041-7.
85	Reaction of 9,10-dimethylacridinium chloride with methanol. Nucleophilic addition of methanol and spontaneous heterolysis of the methanol adduct. Kano, K.; Zhou, B.; Hashimoto, S.; Sugino, T.; Torii, A., <i>Chem. Express</i> 1987, 2 (1), 13-16.
86	Thermal and Photochemical Reactions of 10-Methyl-9-phenylacridinium Chloride in Alcohols. Zhou, B.; Kano, K.; Hashimoto, S., <i>Bull. Chem. Soc. Jpn.</i> 1988, 61, 1041-1047.
87	Binding sites of pyrene and related compounds and chiral excimer formation in the cavities of cyclodextrins and branched cyclodextrins. Kano, K.; Matsumoto, H.; Yoshimura, Y.; Hashimoto, S., <i>J. Am. Chem. Soc.</i> 1988, 110 (1), 204-9.
88	Enantioselective complexation of bilirubin with cyclodextrins and noncyclic oligosaccharides. Kano, K.; Yoshiyasu, K.; Hashimoto, S., <i>J. Chem. Soc., Chem. Commun.</i> 1988, (12), 801-2.
89	Nonosecond Pulse Fluorometry of Conformational Change in Phenylalanine Hydroxylase Associated with Activation. Koizumi, S.; Tanaka, F.; Kaneda, N.; Kano, K.; Nagatsu, T., <i>Biochemistry</i> 1988, 27, 640-646.
90	Hydride transfer from alcohols to protonated azobenzene. Kano, K.; Tatsumi, M.; Hashimoto, S., <i>Sci. Eng. Rev. Doshisha Univ.</i> 1989, 30 (2), 117-21.
91	Chiral recognition of binaphthyls by permethylated β -cyclodextrin. Kano, K.; Yoshiyasu, K.; Hashimoto, S., <i>J. Chem. Soc., Chem. Commun.</i> 1989, (17), 1278-9.
92	Monolayers of 1-Stearylpyrene-6-sulfonate and 1-Stearylpyrene-8-sulfonate. Kano, K.; Fujii, H.; Uraki, H.; Hashimoto, S., <i>Langmuire</i> 1989, 5, 927-932.
93	Selectivities in cyclodextrin chemistry. Kano, K., <i>Bioorg. Chem. Front.</i> 1993, 3, 1-23.
94	Molecular complex of a cationic porphyrin and p-nitrophenol. Verification of the porphyrin dimerization. Kano, K.; Hashimoto, S., <i>Bull. Chem. Soc. Jpn.</i> 1990, 63 (2), 633-5.
95	Cationic porphyrins in water: proton NMR and fluorescence studies on dimer and molecular complex formation. Kano, K.; Takei, M.; Hashimoto, S., <i>J. Phys. Chem.</i> 1990, 94 (5), 2181-7.
96	Hemin-catalyzed addition reactions of thiophenols to styrene. Kano, K.; Takeuchi, M.; Hashimoto, S.; Yoshida, Z., <i>Chem. Lett.</i> 1990, (8), 1381-4.
97	Molecular recognition by saccharides. Asymmetric complexation between bilirubin and nucleosides. Kano, K.; Yoshiyasu, K.; Hashimoto, S., <i>Chem. Lett.</i> 1990, (1), 21-4.

加納航治
2014年1月末日までの全論文

98	Strong van der Waals interactions in water. Molecular complexes of porphyrins and quinones in organic and aqueous media. Kano, K.; Hayakawa, T.; Hashimoto, S., <i>Bull. Chem. Soc. Jpn.</i> 1991, 64
99	Control of headgroup orientation of monolayer by glucolipid. Kano, K.; Ishimura, T.; Hashimoto, S., <i>J. Phys. Chem.</i> 1991, 95 (20), 7839-43.
100	Porphinatoiron-catalyzed oxygenation of styrene in aqueous solution. Kano, K.; Takagi, H.; Takeuchi, M.; Hashimoto, S.; Yoshida, Z., <i>Chem. Lett.</i> 1991, (3), 519-22.
101	Porphyrinatoiron-catalyzed reduction of styrene with sodium borohydride: proposed σ -alkyliron(III) complex as an intermediate. Kano, K.; Takeuchi, M.; Hashimoto, S.; Yoshida, Z., <i>J. Chem. Soc., Chem. Commun.</i> 1991, (24), 1728-9.
102	Specific interactions between sodium deoxycholate and its water-insoluble analogues: mechanisms for premicelle and micelle formation of sodium deoxycholate. Kano, K.; Tatemoto, S.; Hashimoto, S., <i>J. Phys. Chem.</i> 1991, 95 (2), 966-70.
103	Cyclodextrin-induced conformational enantiomerism of dinaphthylmethanes. Kano, K.; Tatsumi, M.; Hashimoto, S., <i>J. Org. Chem.</i> 1991, 56 (23), 6579-85.
104	Binding forces in complexation of p-alkylphenols with β -cyclodextrin and methylated β -cyclodextrins. Kano, K.; Tamiya, Y.; Hashimoto, S., <i>J. Inclusion Phenom. Mol. Recognit. Chem.</i> 1992, 13 (3), 287-93.
105	Mechanisms of steroid-induced conformational enantiomerism of bilirubin in protic solvents. Kano, K.; Tsujino, N.; Kim, M., <i>J. Chem. Soc., Perkin Trans. 2</i> 1992, (10), 1747-52.
106	Chiral recognition by cyclic oligosaccharides. Enantioselective complexation of bilirubin with β -cyclodextrin through hydrogen bonding in water. Kano, K.; Yoshiyasu, K.; Yasuoka, H.; Hata, S.; Hashimoto, S., <i>J. Chem. Soc., Perkin Trans. 2</i> 1992, (8), 1265-9.
107	Fluorescence and circular dichroism studies on molecular and chiral recognition by cyclodextrins. Kano, K., <i>Inclusion Phenom. Mol. Recognit., [Proc. Int. Symp.]</i> , 5th 1990, 243-9.
108	Chiral recognition by cyclic oligosaccharides. Enantioselective complexation of binaphthyl derivatives with cyclodextrins. Kano, K.; Tamiya, Y.; Otsuki, C.; Shimomura, T.; Ohno, T.; Hayashida, O.; Murakami, Y., <i>Supramol. Chem.</i> 1993, 2 (2-3), 137-43.
109	(σ -Alkyl)iron complexes as intermediates in (porphinato)iron-mediated reduction of alkenes and alkynes with sodium borohydride. Takeuchi, M.; Kano, K., <i>Organometallics</i> 1993, 12 (6), 2059-64.
110	Chiral recognition by oligosaccharides. Kano, K., <i>Yukagaku</i> 1994, 43 (10), 868-77.
111	Mechanisms for (porphinato)manganese(III)-catalyzed oxygenation and reduction of styrenes in benzene-ethanol containing sodium borohydride. Takeuchi, M.; Kano, K., <i>Bull. Chem. Soc. Jpn.</i> 1994, 67 (6), 1726-33.
112	(Porphinato)iron-Catalyzed Addition Reactions of Thiols to Alkenes via (σ -Alkyl)iron(II) Complexes. Takeuchi, M.; Shimakoshi, H.; Kano, K., <i>Organometallics</i> 1994, 13 (4), 1208-13.
113	Conformational enantiomerism of bilirubin and pamoic acid induced by protonated amino cyclodextrins. Kano, K.; Arimoto, S.; Ishimura, T., <i>J. Chem. Soc., Perkin Trans. 2</i> 1995, 1661-6.
114	Properties of alkyl β -D-glucoside and alkyl β -D-maltoside micelles. Kano, K.; Ishimura, T., <i>J. Chem. Soc., Perkin Trans. 2</i> 1995, 1655-60.
115	Mechanism for binding to the flexible cavity of permethylated α -cyclodextrin. Kano, K.; Ishimura, T.; Negi, S., <i>J. Inclusion Phenom. Mol. Recognit. Chem.</i> 1995, 22 (4), 285-98.
116	Ability of non-cyclic oligosaccharides to form molecular complexes and its use for chiral separation by capillary zone electrophoresis. Kano, K.; Minami, K.; Horiguchi, K.; Ishimura, T.; Koderu, M., <i>J. Chromatogr. A</i> 1995, 694, 307-13.
117	Mechanism for chiral recognition of binaphthyl derivatives by cyclodextrins. Kano, K.; Kato, Y.; Koderu, M., <i>J. Chem. Soc., Perkin Trans. 2</i> 1996, (6), 1211-1217.
118	Use of electrostatic interaction for chiral recognition. Enantioselective complexation of anionic binaphthyls with protonated amino- β -cyclodextrin. Kano, K.; Kitae, T.; Takashima, H., <i>J. Inclusion Phenom. Mol. Recognit. Chem.</i> 1996, 25 (1-3), 243-248.
119	Preparation of poly(vinyl chloride)/amphiphilic cyclodextrin composite films and their chiral recognition ability. Kano, K.; Takashima, H.; Taniuchi, Y., <i>Kobunshi Ronbunshu</i> 1996, 53, 542-547.
120	Tetraarylporphyrins as probes for studying mechanism of inclusion-complex formation of cyclodextrins. Effects of microscopic environment on inclusion of ionic guests. Kano, K.; Tanaka, N.; Minamizono, H.; Kawakita, Y., <i>Chem. Lett.</i> 1996, 925-926.
121	First example of a rigid (μ -oxo-di- μ -acetato)diiron(III) complex with 1,2-bis[2-di-2-pyridylmethyl-6-pyridyl]ethane; its efficient catalysis for functionalization of alkanes. Koderu, M.; Shimakoshi, H.; Kano, K., <i>Chem. Commun. (Cambridge)</i> 1996, (15), 1737-1738.

加納航治
2014年1月末日までの全論文

122	Synthesis, Characterization, and Crystal Structure of a (μ -Oxo)bis(μ -acetato)diiron(III) Complex with a Dinucleating Hexapyridine Ligand, 1,2-Bis[2-(bis(2-pyridyl)methyl)-6-pyridyl]ethane. The First Example of a Discrete (μ -Oxo)bis(μ -acetato)diiron(III) Complex with a Dinucleating Ligand. Kodera, M.; Shimakoshi, H.; Nishimura, M.; Okawa, H.; Iijima, S.; Kano, K., <i>Inorg. Chem.</i> 1996, 35,
123	Mechanisms for (porphyrinato)iron(III)-catalyzed oxygenation of styrenes by O ₂ in presence of BH ₄ . Takeuchi, M.; Kodera, M.; Kano, K.; Yoshida, Z.-i., <i>J. Mol. Catal. A: Chem.</i> 1996, 113, 51-59.
124	Mechanisms for chiral recognition by cyclodextrins. Kano, K., <i>J. Phys. Org. Chem.</i> 1997, 10, 286-
125	Entropy-dominating strong binding of carboxylate anions to protonated aminocyclodextrin. Kano, K.; Kitae, T.; Takashima, H.; Shimofuri, Y., <i>Chem. Lett.</i> 1997, 899-900.
126	Self-Aggregation of Cationic Porphyrins in Water. Can π - π Stacking Interaction Overcome Electrostatic Repulsive Force? Kano, K.; Minamizono, H.; Kitae, T.; Negi, S., <i>J. Phys. Chem. A</i> 1997, 101, 6118-6124.
127	Optical resolution of 1-arylethanol using transesterification catalyzed by lipases. Kano, K.; Negi, S.; Kawashima, A.; Nakamura, K., <i>Enantiomer</i> 1997, 2, 261-266.
128	Chiral recognition of tetrahelicene dicarboxylic acid by linear dextrans. Kano, K.; Negi, S.; Takaoka, R.; Kamo, H.; Kitae, T.; Yamaguchi, M.; Okubo, H.; Hirma, M., <i>Chem. Lett.</i> 1997, 715-716.
129	Synthesis, Crystal Structures, and Catalytic Activity of Dicopper(II) Complexes with Dinucleating Tetraimidazole Ligands [Cu ₂ (RO)(mbipl)](ClO ₄) ₂ (R = Me, Et, 2-Pr; Hmbipl = 1,5-Bis[bis[(1-methyl-4-imidazolyl)methyl]amino]-3-pentanol) and [Cu ₂ (MeO)(pbipl)](ClO ₄) ₂ (Hpbipl = 1,5-Bis[bis[(1-isopropyl-4-imidazolyl)methyl]amino]-3-pentanol). Kodera, M.; Terasako, N.; Kita, T.; Tachi, Y.; Kano, K.; Yamazaki, M.; Koikawa, M.; Tokii, T., <i>Inorg. Chem.</i> 1997, 36, 3861-3868.
130	Recognition of helicity by native cyclodextrins. Highly enantioselective complexation of tetrahelicene dicarboxylic acid with β -cyclodextrin. Kano, K.; Negi, S.; Kamo, H.; Kitae, T.; Yamaguchi, M.; Okubo, H.; Hirma, M., <i>Chem. Lett.</i> 1998, 151-152.
131	Chiral recognition of α -amino acids by charged cyclodextrins through cooperative effects of Coulomb interaction and inclusion. Kitae, T.; Nakayama, T.; Kano, K., <i>J. Chem. Soc., Perkin Trans. 2</i> 1998,, 207-212.
132	Efficient oxidation of various phenols catalyzed by Di- μ -hydroxodicopper(II) complexes of a hexapyridine dinucleating ligand. Kodera, M.; Shimakoshi, H.; Tachi, Y.; Katayama, K.; Kano, K., <i>Chem. Lett.</i> 1998, (5), 441-442.
133	A superoxodicopper(II) complex oxidatively generated by a reaction of di- μ -hydroxodicopper(II) complex with hydrogen peroxide. Kodera, M.; Tachi, Y.; Hirota, S.; Katayama, K.; Shimakoshi, H.; Kano, K.; Fujisawa, K.; Moro-Oka, Y.; Naruta, Y.; Kitagawa, T., <i>Chem. Lett.</i> 1998, 389-390.
134	Chiral recognition of 1-(4-quinolyl)ethanol by permethylated α -cyclodextrin. Negi, S.; Terai, K.; Kano, K.; Nakamura, K., <i>J. Chem. Res., Synop.</i> 1998, 750-751, 3268-3296.
135	Chiral recognition of an anionic tetrahelicene by native cyclodextrins. Enantioselectivity dominated by location of a hydrophilic group of the guest in a cyclodextrin cavity. Kano, K.; Kamo, H.; Negi, S.; Kitae, T.; Takaoka, R.; Yamaguchi, M.; Okubo, H.; Hirma, M., <i>J. Chem. Soc., Perkin Trans. 2</i> 1999,
136	Recognition of planar chirality by cyclodextrins. Kano, K.; Takaoka, R.; Sato, M.; Yamaguchi, M., <i>Chem. Lett.</i> 1999, 1337-1338.
137	Chiral recognition of phenylacetic acid derivatives by aminated cyclodextrins. Kitae, T.; Takashima, H.; Kano, K., <i>J. Inclusion Phenom. Macrocyclic Chem.</i> 1999, 33, 345-359.
138	Crystal Structure and Reversible O ₂ -Binding of a Room Temperature Stable micro- η^2 : η^2 -Peroxodicopper(II) Complex of a Sterically Hindered Hexapyridine Dinucleating Ligand. Kodera, M.; Katayama, K.; Tachi, Y.; Kano, K.; Hirota, S.; Fujinami, S.; Suzuki, M., <i>J. Am. Chem. Soc.</i> 1999, 121 (47), 11006-11007.
139	Factors Influencing Self-Aggregation Tendencies of Cationic Porphyrins in Aqueous Solution. Kano, K.; Fukuda, K.; Wakami, H.; Nishiyabu, R.; Pasternack, R. F., <i>J. Am. Chem. Soc.</i> 2000, 122, 7494-
140	Complexation of polyvalent cyclodextrin ions with oppositely charged guests: entropically favorable complexation due to dehydration. Kano, K.; Kitae, T.; Shimofuri, Y.; Tanaka, N.; Mineta, Y., <i>Chem. - Eur. J.</i> 2000, 6, 2705-2713.
141	A Cu(II)-Mediated C-H Oxygenation of Sterically Hindered Tripyridine Ligands To Form Triangular Cu(II) ₃ Complexes. Kodera, M.; Tachi, Y.; Kita, T.; Kobushi, H.; Sumi, Y.; Kano, K.; Shiro, M.; Koikawa, M.; Tokii, T.; Ohba, M.; Okawa, H., <i>Inorg. Chem.</i> 2000, 39, 226-234.
142	Effects of Anions in Subphases on a Langmuir Monolayer of N,N-Dimethyl-1-octadecanamine. Negi, S.; Futami, E.; Tsukube, H.; Kano, K., <i>Bull. Chem. Soc. Jpn.</i> 2000, 73, 977-984.
143	Optical resolution of (+)-1-aryl-1-alkanols using enantioselective transesterification by lipases. Negi, S.; Umetsu, K.; Nishijo, Y.; Kano, K.; Nakamura, K., <i>Enantiomer</i> 2000, 5 (1), 63-70.

加納航治
2014年1月末日までの全論文

144	Chiral Recognition of Ru(phen) ₃ ²⁺ by Anionic Cyclodextrins. Kano, K.; Hasegawa, H. <i>Chem. Lett.</i> 2000, 29, 698–699.
145	Interactions with charged cyclodextrins and chiral recognition. Kano, K.; Hasegawa, H., <i>J. Inclusion Phenom. Macrocyclic Chem.</i> 2001, 4, 41–47.
146	Chiral recognition of dipeptide methyl esters by an anionic β-cyclodextrin. Kano, K.; Hasegawa, H.; Miyamura, M., <i>Chirality</i> 2001, 13 (8), 474–482.
147	Hydrogen-bonded complexes of carboxylate anions and dextrins in an aprotic polar solvent. Kano, K.; Tanaka, N.; Negi, S., <i>Eur. J. Org. Chem.</i> 2001, 3689–3694.
148	Hydroperoxo-Copper(II) Complex Stabilized by N ₃ S-Type Ligand Having a Phenyl Thioether. Kodera, M.; Kita, T.; Miura, I.; Nakayama, N.; Kawata, T.; Kano, K.; Hirota, S., <i>J. Am. Chem. Soc.</i> 2001, 123, 7715–7716.
149	Synthesis, Characterization, and Activation of Thermally Stable μ-1,2-Peroxodiiron(III) Complex. Kodera, M.; Taniike, Y.; Itoh, M.; Tanahashi, Y.; Shimakoshi, H.; Kano, K.; Hirota, S.; Iijima, S.; Ohba, M.; Okawa, H., <i>Inorg. Chem.</i> 2001, 40 (19), 4821–4822.
150	Static and Dynamic Behavior of 2:1 Inclusion Complexes of Cyclodextrins and Charged Porphyrins in Aqueous Organic Media. Kano, K.; Nishiyabu, R.; Asada, T.; Kuroda, Y., <i>J. Am. Chem. Soc.</i> 2002, 124, 9937–9944.
151	General mechanism for chiral recognition by native and modified cyclodextrins. Kano, K.; Nishiyabu, R., <i>J. Inclusion Phenom. Macrocyclic Chem.</i> 2002, 44, 355–359.
152	Reexamination of cyclodextrin-induced conformational enantiomerism of bilirubin in aqueous solution. Kano, K.; Imaeda, K.; Ota, K.; Doi, R., <i>Bull. Chem. Soc. Jpn.</i> 2003, 7, 1035–1041.
153	Polar effect on the stacking interaction between a zinc complex of an anionic porphyrin and anionic aromatics. Kano, K.; Kobayashi, S., <i>Bull. Chem. Soc. Jpn.</i> 2003, 7, 2027–2034.
154	Chiral recognition by cyclodextrins: a general mechanism. Kano, K.; Nishiyabu, R., <i>Adv. Supramol. Chem.</i> 2003, 9, 39–69.
155	Convenient Scaffold for Forming Heteroporphyrin Arrays in Aqueous Media. Kano, K.; Nishiyabu, R.; Yamazaki, T.; Yamazaki, I., <i>J. Am. Chem. Soc.</i> 2003, 125, 10625–10634.
156	Structural Modulation of Cu(I) and Cu(II) Complexes of Sterically Hindered Tripyridine Ligands by the Bridgehead Alkyl Groups. Kodera, M.; Kajita, Y.; Tachi, Y.; Kano, K., <i>Inorg. Chem.</i> 2003, 42,
157	Mechanism for aerobic oxidation of 3,5-di-tert-butylcatechol to 3,5-di-tert-butyl-o-benzoquinone catalyzed by di-μ ⁻ hydroxo-dicopper(II) complexes of peralkylated ethylenediamine ligands. Kodera, M.; Kawata, T.; Kano, K.; Tachi, Y.; Itoh, S.; Kojo, S., <i>Bull. Chem. Soc. Jpn.</i> 2003, 76, 1957–
158	Molecular complexes of water-soluble porphyrins. Kano, K., <i>J. Porphyrins Phthalocyanines</i> 2004, 8, 148–155.
159	Heptakis(2,3-di-O-carboxymethyl)-β-cyclodextrin as a pH-sensitive host. Kano, K.; Horiki, Y.; Mabuchi, T.; Kitagishi, H., <i>Chem. Lett.</i> 2004, 33, 1086–1087.
160	Anion Binding to a Ferric Porphyrin Complexed with Per-O-methylated β-Cyclodextrin in Aqueous Solution. Kano, K.; Kitagishi, H.; Tamura, S.; Yamada, A., <i>J. Am. Chem. Soc.</i> 2004, 126 (46), 15202–15210.
161	Synthesis, structure, and greatly improved reversible O ₂ binding in a structurally modulated μ-η ² : η ² -peroxodicopper(II) complex with room-temperature stability. Kodera, M.; Kajita, Y.; Tachi, Y.; Katayama, K.; Kano, K.; Hirota, S.; Fujiwara, S.; Suzuki, M., <i>Angew. Chem., Int. Ed.</i> 2004, 43, 334–
162	Double self-inclusion by rotating glucopyranose units in per-O-methylated β-cyclodextrin moieties attached to a porphyrin in aqueous solution. Nishiyabu, R.; Kano, K., <i>Eur. J. Org. Chem.</i> 2004, (24), 4985–4988.
163	Copper(II) Complexes of a Series of Polypyridine Ligands Possessing a 1,2-Bis(2-pyridyl)ethane Common Moiety: Incorporation and Hydrolysis of Phosphate Esters. Itoh, M.; Nakazawa, J.; Maeda, K.; Kano, K.; Mizutani, T.; Kodera, M., <i>Inorg. Chem.</i> 2005, 44, 691–702.
164	Dioxygen binding to a simple myoglobin model in aqueous solution. Kano, K.; Kitagishi, H.; Kodera, M.; Hirota, S., <i>Angew. Chem., Int. Ed.</i> 2005, 44, 435–438.
165	Novel Behavior of O-Methylated β-Cyclodextrins in Inclusion of meso-tetra-Arylporphyrins. Kano, K.; Nishiyabu, R.; Doi, R., <i>J. Org. Chem.</i> 2005, 70, 3667–3673.
166	A diiron center stabilized by a bis-TPA ligand as a model of soluble methane monooxygenase: Predominant alkene epoxidation with H ₂ O ₂ . Kodera, M.; Itoh, M.; Kano, K.; Funabiki, T.; Reglier, M., <i>Angew. Chem., Int. Ed.</i> 2005, 44, 7104–7106.
167	Iron Porphyrin-Cyclodextrin Supramolecular Complex as a Functional Model of Myoglobin in Aqueous Solution. Kano, K.; Kitagishi, H.; Dagallier, C.; Kodera, M.; Matsuo, T.; Hayashi, T.; Hisaeda, Y.; Hirota, S., <i>Inorg. Chem.</i> 2006, 45, 4448–4460.

加納航治
2014年1月末日までの全論文

168	A myoglobin functional model composed of a ferrous porphyrin and a cyclodextrin dimer with an imidazole linker. Kano, K.; Kitagishi, H.; Mabuchi, T.; Kodera, M.; Hirota, S., <i>Chem. – Asian J.</i> 2006, 1, 358–366.
169	Redox behavior of a manganese porphyrin complexed with per-O-methylated β -cyclodextrin in aqueous solution. Kano, K.; Kitagishi, H.; Sone, Y.; Nakazawa, N.; Kodera, M., <i>Eur. J. Inorg. Chem.</i> 2006, 4043–4053.
170	Characterization of an Inclusion Complex of 5,10,15,20-Tetrakis(4-sulfonatophenyl)porphyrinato iron and an O-Methylated β -Cyclodextrin Dimer Having a Pyridine Linker and Its Related Complexes in Aqueous Solution. Kano, K.; Kitagishi, H.; Tanaka, S., <i>J. Inclusion Phenom. Macrocylic Chem.</i> 2006, 56, 69–74.
171	Peroxodiiron complexes of polypyridine ligands: syntheses, physicochemical properties, and thermal stability markedly enhanced by hexapyridine dinucleating ligand. Kodera, M.; Itoh, M.; Kano, K.; Funabiki, T., <i>Bull. Chem. Soc. Jpn.</i> 2006, 7, 252–261.
172	Thermodynamics of hydrophobic interactions: entropic recognition of a hydrophobic moiety by poly(ethylene oxide)-zinc porphyrin conjugates. Iwamoto, H.; Mizutani, T.; Kano, K., <i>Chem. – Asian J.</i> 2007, 2, 1267–1275.
173	Supramolecular complex of cytochrome c with a polyanionic β -cyclodextrin. Kano, K.; Ishida, Y., <i>Angew. Chem., Int. Ed.</i> 2007, 46, 727–730, S727/1–S727/10.
174	Heat-capacity changes in host-guest complexation by Coulomb interactions in aqueous solution. Kano, K.; Ishida, Y.; Kitagawa, K.; Yasuda, M.; Watanabe, M., <i>Chem. – Asian J.</i> 2007, 2, 1305–1313.
175	Reversible O ₂ -binding and activation with dicopper and diiron complexes stabilized by various hexapyridine ligands. stability, modulation, and flexibility of the dinuclear structure as key aspects for the dimetal/O ₂ chemistry. Kodera, M.; Kano, K., <i>Bull. Chem. Soc. Jpn.</i> 2007, 80, 662–676.
176	Porphyrin-cyclodextrin supramolecular complexes as myoglobin model in water. Kano, K., <i>Colloid Polym. Sci.</i> 2008, 286, 79–84.
177	Regulation of α -chymotrypsin catalysis by ferric porphyrins and cyclodextrins. Kano, K.; Ishida, Y., <i>Chem. – Asian J.</i> 2008, 3, 678–686.
178	A Supramolecular Receptor of Diatomic Molecules (O ₂ , CO, NO) in Aqueous Solution. Kano, K.; Itoh, Y.; Kitagishi, H.; Hayashi, T.; Hirota, S., <i>J. Am. Chem. Soc.</i> 2008, 130, 8006–8015.
179	Porphyrin J-Aggregates Stabilized by Ferric Myoglobin in Neutral Aqueous Solution. Kano, K.; Watanabe, K.; Ishida, Y., <i>J. Phys. Chem. B</i> 2008, 112, 14402–14408.
180	HemoCD as an artificial oxygen carrier: oxygen binding and autoxidation. Kano, K.; Kitagishi, H., <i>Artif. Organs</i> 2009, 33, 177–182.
181	Novel zinc finger nuclease created by combining the Cys ₂ His ₂ and His ₄ type zinc finger domains. Negi, S.; Umeda, Y.; Masuyama, S.; Kano, K.; Sugiura, Y., <i>Bioorg. Med. Chem. Lett.</i> 2009, 19, 2789–2791.
182	A diatomic molecule receptor that removes CO in a living organism. Kitagishi, H.; Negi, S.; Kiriya, A.; Honbo, A.; Sugiura, Y.; Kawaguchi Akira, T.; Kano, K., <i>Angew. Chem., Int. Ed. Engl.</i> 2010, 49 (7),
183	Oxoferryl Porphyrin/Hydrogen Peroxide System Whose Behavior is Equivalent to Hydroperoxoferric Porphyrin. Kitagishi, H.; Tamaki, M.; Ueda, T.; Hirota, S.; Ohta, T.; Naruta, Y.; Kano, K., <i>J. Am. Chem. Soc.</i> 2010, 13, 16730–16732.
184	Time-Dependent Enzyme Activity Dominated by Dissociation of J-Aggregates Bound to Protein Surface. Watanabe, K.; Kano, K., <i>Bioconjugate Chem.</i> 2010, 2, 2332–2338.
185	Binding of Multivalent Anionic Porphyrins to V3 Loop Fragments of an HIV-1 Envelope and Their Antiviral Activity. Watanabe, K.; Negi, S.; Sugiura, Y.; Kiriya, A.; Honbo, A.; Iga, K.; Kodama, E. N.; Naitoh, T.; Matsuoka, M.; Kano, K., <i>Chem. – Asian J.</i> 2010, 5, 825–834.
186	Structural Basis of the Lactate-dependent Allosteric Regulation of Oxygen Binding in Arthropod Hemocyanin. Hirota, S.; Tanaka, N.; Micetic, I.; Di Muro, P.; Nagao, S.; Kitagishi, H.; Kano, K.; Magliozzo, R. S.; Pelsach, J.; Beltramini, M.; Bubacco, L. <i>J. Biol. Chem.</i> 2010, 285(25), 19338–19345.
187	Completely synthetic compounds that bind diatomic molecules (O ₂ , CO, NO) in aqueous solution. Kitagishi, H.; Kano, K., <i>J. Inorg. Nucl. Chem.</i> 2010, 17, 160–168.
188	Calorimetric Study on Coordination of Tridentate Imidazolyl Calix[6]arene Ligands to Zinc Ion in Organic Solvents. Kano, K.; Kondo, M.; Inoue, H.; Kitagishi, H.; Colasson, B.; Reinaud, O., <i>Inorg. Chem.</i> 2011, 50, 6353–6360.
189	Preparation and Function of Poly(acrylic acid)s Modified by Supramolecular Complex Composed of Porphyrinatoiron and a Cyclodextrin Dimer That Bind Diatomic Molecules (O ₂ and CO) in Aqueous Solution. Kano, K.; Ochi, T.; Okunaka, S.; Ota, Y.; Karasugi, K.; Ueda, T.; Kitagishi, H., <i>Chem. – Asian J.</i> 2011, 6, 2946–2955.

加納航治
2014年1月末日までの全論文

190	Gold Nanoparticles Carrying Diatomic Molecules (O ₂ and CO) in Aqueous Solution. Karasugi, K.; Kitagishi, H.; Kano, K., Chem. – Asian J. 2011, 6, 825–833.
191	Supramolecular Ferric Porphyrins as Cyanide Receptors in Aqueous Solution. Watanabe, K.; Kitagishi, H.; Kano, K., ACS Med. Chem. Lett. 2011, 2, 943–947.
192	Supramolecular dioxygen receptors composed of an anionic water-soluble porphyrinatoiron(II) and cyclodextrin dimers. Kano, K.; Chimoto, S.; Tamaki, M.; Itoh, Y.; Kitagishi, H., Dalton Trans. 2012, 41, 453–461.
193	Modification of a Dioxygen Carrier, HemoCD, with PEGylated Dendrons for Extension of Circulation Time in the Bloodstream. Karasugi, K.; Kitagishi, H.; Kano, K., Bioconjugate Chem. 2012, 23, 2365–
194	Functionalization of a protein surface with per-O-methylated β -cyclodextrin. Kitagishi, H.; Kashiwa, K.; Kano, K., Biopolymers 2012, 97, 11–20.
195	PEGylation of an artificial O ₂ and CO receptor: synthesis, characterisation and pharmacokinetic study. Ueda, T.; Kitagishi, H.; Kano, K., Org. Biomol. Chem. 2012, 10, 4337–4347.
196	Elongation of circulation time of a diatomic molecule receptor in bloodstream by attachment to small gold nanoparticles. Karasugi, K.; Kitagishi, H.; Kano, K., Chem. Lett. 2013, 42, 115–117.
197	Cellular uptake of octaarginine-conjugated tetraarylporphyrin included by per-O-methylated β -cyclodextrin. Kitagishi, H.; Hatada, S.; Itakura, T.; Maki, Y.; Maeda, Y.; Kano, K., Org. Biomol. Chem. 2013, 11, 3203–3211.
198	Supramolecular Iron Porphyrin/Cyclodextrin Dimer Complex that Mimics the Functions of Hemoglobin and Methemoglobin. Watanabe, K.; Kitagishi, H.; Kano, K., Angew. Chem., Int. Ed. 2013,
199	Intramolecular Oxidative O-Demethylation in a Per-O-methylated β-Cyclodextrin-Iron Porphyrin Inclusion Complex in Aqueous Solution. Ueda, T.; Kumeda, S.; Kitagishi, H.; Kano, K. Chem. Lett. 2013, 42(11), 1366–1368.
200	Intramolecular Direct Oxygen Transfer from Oxoferryl Porphyrin to a Sulfide Bond. Ueda, T.; Kitagishi, H.; Kano, K. Inorg. Chem. 2014, 53 (1), 543–551.